


Section A

1.

(b) Zn – Hg with HCl

Explanation:



This is called as clemmensen reduction.

2.

(b) All of these

Explanation:

When a haloalkane with β -hydrogen atom is heated with an alcoholic solution of potassium hydroxide, there is an elimination of hydrogen atom from β -carbon and a halogen atom from the α -carbon atom. As a result, an alkene is formed as a product. Since the β -hydrogen atom is involved in elimination, it is often called β -elimination.

3.

(a) Proline

Explanation:

Those amino acids which cannot be synthesised in the body and must be obtained through diet, are known as essential amino acids. Proline is not an essential amino acid.

4.

(d) I_2 and Na_2CO_3

Explanation:



Acetophenone contain $\text{CH}_3 - \text{C}(=\text{O}) -$ unit thus it will give haloform test and precipitate CHI_3 (yellow ppt). On other hand

benzophenone does not form haloform.

5.

(c) All of these

Explanation:

Phenol reacts with sodium hydroxide solution to give a colourless solution containing sodium phenoxide. In this reaction, the hydrogen ion has been removed by the strongly basic hydroxide ion in the sodium hydroxide solution.

6.

(d) (a) - (ii), (b) - (iv), (c) - (i), (d) - (iii)

Explanation:

(a) - (ii), (b) - (iv), (c) - (i), (d) - (iii)

7.

(b) four times

Explanation:

four times

8. (a) gem-dihalide


Explanation:

Gem-dihalides are dihaloalkanes that have two halogen atoms of the same type attached to the same carbon atom in a molecule.


The common naming system of gem-dihalides (geminal halide) is alkylidene dihalides. Ethylidene dichloride thus is a gem-

dihalide. The chemical formula of ethylidene dichloride is $C_3H_6Cl_2$.

9.




Explanation:



10.

(b)



Explanation:

According to Arrhenius equation $k = Ae^{-E_a/RT}$

Taking log on both sides in $k = \ln \left(A \cdot e^{-\frac{E_a}{RT}} \right)$

$$\ln k = \ln A - \frac{E_a}{RT}$$

$$\ln k = -\frac{E_a}{R} \times \frac{1}{T} + \ln A$$

$$y = mx + c$$

This equation can be related to the equation of a straight line.

From the graph, it is very clearly shown that the slope of the plot $= \frac{-E_a}{R}$ and intercept $= \ln A$.

11. (a) Tertiary alcohol

Explanation:

The outcome of oxidation reactions of alcohol depends on the substituents on the carbinol carbon. In order for each oxidation step to occur, there must be H on the carbinol carbon.

- Primary alcohols can be oxidized to aldehydes or further to carboxylic acids. In aqueous media, the carboxylic acid is usually the major product. PCC or PDC, which are used in dichloromethane, allow the oxidation to be stopped at the intermediate aldehyde.
- Secondary alcohols can be oxidized to ketones but no further:
- Tertiary alcohols cannot be oxidized (no carbinol C-H).

12.

(c) They are chemically reactive.

Explanation:

Interstitial compounds are obtained when small atoms like H, B, C, resemble N, etc. fit into the lattice of other elements. They are chemically inert.

13.

(c) A is true but R is false.

Explanation:

A is true but R is false.

14.

(c) A is true but R is false.

Explanation:

$\text{CH}_2=\text{CH}-\text{Cl}$ has some partial double bond character between carbon and a chlorine atom. So, nucleophilic substitution is difficult to carry as it is difficult to break the partial double bond in vinyl chloride than ethyl chloride $\text{CH}_3\text{CH}_2-\text{Cl}$ where there is no double bond character.

The vinyl group is not electron-donating. The carbon halogen bond in vinyl halides has some double-bond character and thus a little difficult to break.

15.

(c) A is true but R is false.

Explanation:

A is true but R is false.

16.

(b) Both A and R are true but R is not the correct explanation of A.

Explanation:

Both A and R are true but R is not the correct explanation of A.

Section B

17. Examples of complexes in biological system.-

1. Chlorophyll is a complex of Mg.
2. Haemoglobin is a complex of iron.
3. Cyanocobalamin, Vitamin B₁₂, is a complex of cobalt.



19. Henry's Law states that "the partial pressure of the gas in vapour phase (p) is proportional to the mole fraction of the gas (x) in the solution" and is expressed as:

$$p = K_H \cdot x$$

Here K_H is the Henry's law constant.

Given, $T = 298 \text{ K}$, $K_H = 1.25 \times 10^6$, $p = 760 \text{ mm Hg}$

We know by Henry's Law, $p = K_H \times x$, where x is mole fraction of the gas in the solution.

$$760 = 1.25 \times 10^6 \times x$$

$$x = \frac{760}{1.25 \times 10^6}$$

$$x = 6.08 \times 10^{-4}$$

OR

Molality: Molality of a solution is defined as the number of moles of the solute dissolved in 1000 g (1 Kg) of the solvent. It is denoted as 'm'.

$$\text{Molality (m)} = \frac{\text{Moles of solute}}{\text{Mass of solvent in Kg}} \quad \text{Unit of molality} = \text{mol/kg}$$

For example, 1.00 mol Kg⁻¹ (or 1.00 m) solution of KCl means that 1 mol of KCl dissolved in 1 Kg of water.

20. Answer the following:

- (i) Rate of reaction is defined as change in concentration of reactants or products per unit time. For example, the reaction $\text{A} \rightarrow \text{B}$ has the rate expressed as:


$$\text{rate of reaction} = \frac{dx}{dt} = \frac{-[dA]}{dt} = \frac{[dB]}{dt}$$

(ii) For a reaction $R \rightarrow P$, half-life ($t_{1/2}$) is observed to be independent of the initial concentration of reactants. Thus, it follows first order reaction.

21. a. $2\text{MnO}_2 + 4\text{KOH} + \text{O}_2 \rightarrow 2\text{K}_2\text{MnO}_4 + 2\text{H}_2\text{O}$
 b. $10\text{I}^- + 2\text{MnO}_4^- + 16\text{H}^+ \rightarrow 2\text{Mn}^{2+} + 8\text{H}_2\text{O} + 5\text{I}_2$
 c. $\text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+ + 3\text{Sn}^{2+} \rightarrow 2\text{Cr}^{3+} + 3\text{Sn}^{4+} + 7\text{H}_2\text{O}$

Above mentioned reactions are complete reactions of given reagents.

Section C



23. i. Average rate of reaction between the time interval, 30 to 60 seconds, $= \frac{d[\text{Ester}]}{dt}$
 $= \frac{0.31-0.17}{60-30} = \frac{0.14}{30} = 4.67 \times 10^{-3} \text{ mol L}^{-1} \text{ s}^{-1}$

ii. For a pseudo first order reaction,

$$k = \frac{2.303}{t} \log \frac{[R]_0}{[R]}$$

$$\text{For } t = 30\text{s}, k_1 = \frac{2.303}{30} \log \frac{0.55}{0.31} = 1.911 \times 10^{-2} \text{ s}^{-1}$$




$$\text{For } t = 60\text{s}, k_2 = \frac{2.303}{60} \log \frac{0.55}{0.17} = 1.957 \times 10^{-2} \text{ s}^{-1}$$

For $t = 90\text{s}$,

$$k_3 = \frac{2.303}{90} \log \frac{0.55}{0.085} = 2.075 \times 10^{-2} \text{ s}^{-1}$$

$$\text{Then, average rate constant, } k = \frac{k_1 + k_2 + k_3}{3} = 1.981 \times 10^{-2} \text{ s}^{-1}$$

24. a. The movement of colloidal particles under an applied electric potential.
 b. Yes this process can be used in the coagulation of the lyophobic sols.
 c. The process of settling colloidal particles is coagulation.



$$E_{\text{cell}} = E_{\text{cell}}^0 - \frac{0.0591}{2} \log \frac{[\text{Mg}^{2+}]}{[\text{Cu}^{2+}]}$$

$$E_{\text{cell}}^0 = \left[E^0 \frac{(\text{Cu}^{2+})}{(\text{Cu})} - E^0 \frac{(\text{Mg}^{2+})}{(\text{Mg})} \right]$$

$$E_{\text{cell}} = [+0.34\text{V} - (-2.37\text{V})] - \frac{0.0591}{2} \log 10^2$$

$$= (0.271\text{V} - 0.0591\text{V})$$



$$= 2.65\text{V}$$

26. The series of elements which have been arranged on the basis of their electrode potential is called electrochemical series or activity series.

Standard EMF of the cell = [standard reduction potential of the right hand side electrode] - [Standard reduction potential of the left hand side electrode]


$$emf = E_{\text{cathode}}^0 - E_{\text{anode}}^0$$

27. This is cross aldol condensation reaction.




28. The mechanism of the reaction of HI with methoxymethane involves the following steps:


Step1: Protonation of methoxymethane:



Step2: Nucleophilic attack of I^- :



Step3: When HI is in excess and the reaction is carried out at a high temperature, the methanol formed in the second step reacts with another HI molecule and gets converted to methyl iodide




OR

- i. KMnO_4/KOH (alkaline KMnO_4)
- ii. $\text{Cu}/573 \text{ K}$ (Hot reduced copper)
- iii. $\text{Br}_2(\text{aq})$ (Bromine water)

Section D

29. Read the text carefully and answer the questions:

The f-block consists of the two series, lanthanoids (the fourteen elements following lanthanum) and actinoids (the fourteen elements following actinium). Because lanthanum closely resembles the lanthanoids. The chemistry of the actinoids is much more complicated. The complication arises partly owing to the occurrence of a wide range of oxidation states in these elements and partly because their radioactivity creates special problems in their study. The overall decrease in atomic and ionic radii from lanthanum to lutetium (the lanthanoid contraction) is a unique feature in the chemistry of the lanthanoids. In the lanthanoids, La(II) and Ln(III) compounds are predominant species.



- (i) Copper exhibits +1 oxidation state more frequently i.e., Cu^{+1} because of its electronic configuration $3d^{10}4s^1$. It can easily lose $4s^1$ electron to give stable $3d^{10}$ configuration.

OR

Because of stronger metallic bonding and high enthalpies of atomization.

- (ii) The ability of O_2 to stabilize higher oxidation states exceeds that of fluorine because oxygen can form multiple bonds with metals.
- (iii) Due to lanthanoid contraction in second series after lanthanum, the atomic radii of elements of second and third series become almost same and hence show similarities in properties.

30. Read the text carefully and answer the questions:

Many chemical and biological processes depend on osmosis, the selective passage of solvent molecules through the porous membrane from a dilute solution to a more concentrated one. The osmotic pressure π depends on molar concentration of the solution ($\pi = CRT$). If two solutions are of equal solute concentration and, hence, have the same osmotic pressure, they are said to

be isotonic. If two solutions are of unequal osmotic pressures, the more concentrated solution is said to be hypertonic and the more diluted solution is described as hypotonic.

Osmosis is the major mechanism, for transporting water upward in the plants. Transpiration in the leaves supports the transport mechanism of water. The osmotic pressure of seawater is about 30 atm; this is the pressure that must be applied to the seawater (separated from pure water using a semi-permeable membrane) to get drinking water.

- (i) A plant cell gets shrink when it is kept in a hypertonic solution.
- (ii) 1.2% sodium chloride solution is hypertonic with respect to 0.9% sodium chloride solution or blood cells. When blood cells are placed in this solution, water flows out of the cells and they shrink due to loss of water by osmosis.
- (iii) When the external pressure applied becomes more than the osmotic pressure of the solution, then the solvent molecules from the solution pass through the semipermeable membrane to the solvent side. This process is called reverse osmosis.


OR

In an upward direction, osmosis helps in the transportation of water in a plant.

Section E

31. Attempt any five of the following:


- (i) It dissolves blood clots and used in the treatment of heart diseases.
- (ii) Milk, carrot
- (iii) Cyclic compound containing element other than carbon i.e., N, S, O at ring position are called heterocyclic bases.
- (iv) Primary structure of proteins tells about the sequence in which various amino acids are linked with each other.
- (v) Vitamin A
- (vi) Keratin and Myosin
- (vii) D-glucose on reaction with NH_2OH (hydroxylamine) yield glucose oxime.



32. I.
 - i. Aniline gets protonated and is deactivated/Aniline on protonation forms anilinium ion which is meta-directing therefore it give good amount of m-nitroaniline.
 - ii. Due to combination of inductive effect and solvation effect.
 - iii. Ammonolysis of alkyl halide is not a good method to prepare pure primary amine because it forms a mixture of amines that is difficult to separate.




ii.



- i. Aniline is a Lewis base and forms a salt with Lewis acid.

Aniline being a Lewis base reacts with Lewis acid (AlCl_3) to form a salt.



As a result, N acquires a positive charge so, it acts as a strong deactivating group for electrophilic substitution reaction. Thus, aniline does not undergo Friedel-Crafts reaction.

- ii. In aqueous solution, basic nature depends on + I-effect, H-bonding, and steric-effect.

The combined effect shows that $(\text{CH}_3)_2\text{NH}$ is more basic than $(\text{CH}_3)_3\text{N}$ as H-bonding is more in case of $(\text{CH}_3)_2\text{NH}$ than in $(\text{CH}_3)_3\text{N}$, which predominates over the stability due to +I- effect of three $-\text{CH}_3$ groups.

- iii. Large pK_b value means a weak base

In aniline, the lone pair of electrons on N-atom is delocalized over the benzene ring. As a result, electron density on the

nitrogen decreases and electrons are not available for donation. In contrast, in $\text{CH}_3\text{-NH}_2$, +I effect of -CH_3 group increases the electron density on the N-atom. Therefore, aniline is a weaker base than methylamine and hence, its pK_b value is higher than that of methylamine.

33. a. $\text{K}[\text{Cr}(\text{H}_2\text{O})_2](\text{C}_2\text{O}_4)_2 \cdot 3\text{H}_2\text{O}$

The IUPAC name = Potassium diaquadioxalatochromate (III) trihydrate.


The Oxidation state of chromium = 3

Electronic configuration: $3d^3 : t_{2g}^3$

Coordination number of compound = 6

Shape: octahedral

Stereochemistry:



Magnetic moment, $\mu = \sqrt{n(n+2)}$

$$= \sqrt{3(3+2)}$$

$$= \sqrt{15} \sim 3.87 \text{ BM}$$

b. $[\text{Co}(\text{NH}_3)_5\text{Cl}]\text{Cl}_2$

The IUPAC name: Pentaamminechloridocobalt(III) chloride

The oxidation state of Co = +3


Coordination number of compound = 6

Shape: octahedral.

Electronic configuration:

$$d^6 : t_{2g}^6$$

Stereochemistry:



Magnetic Moment = 0

c. $\text{CrCl}_3(\text{py})_3$

The IUPAC name: Trichloridotripyridinechromium (III)


The oxidation state of chromium = +3

Electronic configuration for $d^3 = t_{2g}^3$

Coordination number of compound = 6

Shape: octahedral.

Stereochemistry:



Both isomers are optically active. Therefore, a total of 4 isomers exist. Magnetic moment,

$$\mu = \sqrt{n(n+2)} = \sqrt{3(3+2)}$$

$$= \sqrt{15} \sim 3.87 \text{ BM}$$

d. $\text{Cs}[\text{FeCl}_4]$

The IUPAC name : Caesium tetrachloroferrate (III)

The oxidation state of Fe = +3

Electronic configuration of $d^6 = e_g^2 t_{2g}^4$

Coordination number of compound = 4

Shape: tetrahedral

Stereochemistry: optically inactive

Magnetic moment:

$$\mu = \sqrt{n(n+2)} = \sqrt{5(5+2)}$$

$$= \sqrt{35} \sim 5.92 \text{ BM}$$

e. $\text{K}_4[\text{Mn}(\text{CN})_6]$

The IUPAC name = Potassium hexacyanomanganate (II)

The oxidation state of manganese = +2

Electronic configuration: $d^{5+} : t_{2g}^5$

Coordination number of compound = 6

Shape: octahedral.

Stereochemistry: optically inactive Magnetic moment, $\mu = \sqrt{n(n+2)}$
 $= \sqrt{1(1+2)} = \sqrt{3}$
 $= 1.732$

OR

a. The IUPAC name of $[\text{Pt}(\text{NH}_3)_2\text{Cl}(\text{NO}_2)]$ is Diamminechloridonitrito-N-platinum(II).

b. The IUPAC name of $\text{K}_3[\text{Cr}(\text{C}_2\text{O}_4)_3]$ is Potassium trioxalatochromate(III).

- c. The IUPAC name of $[\text{CoCl}_2(\text{en})_2]\text{Cl}$ is Dichloridobis (ethane-1,2-diamine)cobalt(III) chloride.
- d. The IUPAC name of $[\text{Co}(\text{NH}_3)_5(\text{CO}_3)]\text{Cl}$ is Pentaamminecarbonatocobalt(III) chloride.
- e. The IUPAC name of $\text{Hg}[\text{Co}(\text{SCN})_4]$ is Mercury (I) tetrathiocyanatocobaltate(III).



AI POWERED
LEARNING



CONCEPT
MASTERY



PERFORMANCE
FOCUSED



EXCELLENCE
DRIVEN

Powered by [QualCrypt](#)